

(p, q) -form Kähler Electromagnetism

D. CHERNEY^ℳ, E. LATINI^ℒ, AND A. WALDRON^ℳ

^ℳ*Department of Mathematics*
University of California, Davis CA 95616, USA
 cherney,wally@math.ucdavis.edu

^ℒ*Dipartimento di Fisica,*
Università a di Bologna, via Irnerio 46, I-40126 Bologna, Italy and INFN,
Laboratori Nazionali di Frascati, CP 13, I-00044 Frascati, Italy
 latini@lnf.infn.it

ABSTRACT

We present a gauge invariant generalization of Maxwell's equations and p -form electromagnetism to Kähler spacetimes.

In their simplest form our Kähler Maxwell's equations are given by

$$\omega^{\bar{j}} \left[\partial_i \partial_{\bar{j}} A_{i_1 \dots i_p \bar{j}_1 \dots \bar{j}_q} - p \partial_{[i_1} \partial_{\bar{j}} A_{i|i_2 \dots i_p] \bar{j}_1 \dots \bar{j}_q} - q \partial_i \partial_{[\bar{j}_1} A_{i_1 \dots i_p | \bar{j}] \bar{j}_2 \dots \bar{j}_q} \right. \\ \left. + pq \partial_{[i_1} \partial_{[\bar{j}_1} A_{i|i_2 \dots i_p] \bar{j}] \bar{j}_2 \dots \bar{j}_q} \right] = 0. \quad (1)$$

Here $\omega_{i\bar{j}}$ is the Kähler form and we have raised and lowered indices with the Kähler metric, while the dynamical field $A_{i_1 \dots i_p \bar{j}_1 \dots \bar{j}_q}$ is a (p, q) -form on a Kähler manifold M of any complex dimension n . Therefore we call this system of equations (p, q) -form Kähler electromagnetism. To understand this system and its gauge invariances better we first streamline our notations using some basic Hodge-Lefschetz theory (see, for example [1] for a detailed account and original references).

Firstly we denote the Dolbeault decomposition of the exterior derivative d and the codifferential $d^* = \star d \star$ in the usual way

$$d = \partial + \bar{\partial}, \quad d^* = \delta + \bar{\delta}, \quad (2)$$

subject to the $\mathcal{N} = 4$ supersymmetry algebra

$$\{\partial, \delta\} = \frac{1}{2} \Delta = \{\bar{\partial}, \bar{\delta}\}, \quad (3)$$

where Δ is the form Laplacian. The R -symmetries of this superalgebra include the Hodge-Lefschetz $sl(2)$ algebra generated by $\{\Lambda, H, L\}$. The operator Λ is essentially the contraction by the Kähler form appearing in (1), H returns $p+q-n$ on (p, q) forms on an n -dimensional complex manifold while L is the adjoint of Λ with respect to the Hodge inner product. All these operators may be realized in first quantization as symmetries of a quantum mechanical $\mathcal{N} = 4$ supersymmetric sigma model¹ (see [2] for an elegant account and history of this theory). We then have the algebra

$$\begin{aligned} [H, \Lambda] &= -2\Lambda, & [H, L] &= 2L, & [\Lambda, L] &= H, \\ [\Lambda, \partial] &= \bar{\partial}, & [\Lambda, \bar{\partial}] &= -\partial, & [L, \delta] &= \bar{\partial}, & [L, \bar{\delta}] &= -\partial. \end{aligned} \quad (4)$$

In these terms our (p, q) -form Kähler Maxwell's equations become simply

$$\boxed{\Lambda \partial \bar{\partial} A = 0}, \quad (5)$$

for any differential form A on a Kähler manifold M (it is no longer necessary to specify a definite degree for A).

These equations share some remarkable similarities with both Maxwell's equations and the linearized Einstein equations. To see this we must briefly recall the theory of symmetric forms. A symmetric form is any symmetric tensor field compactly expressed as a function of commuting differentials dx^μ (for example the metric tensor $ds^2 = dx^\mu g_{\mu\nu} dx^\nu$). On symmetric forms one has useful geometric operations such as the symmetrized gradient **grad**, divergence **div** and trace **tr** (see [3] for details). In these notations, in a flat background, many physical systems are generated by the equations

$$\left[\square - \mathbf{grad} \mathbf{div} + \frac{1}{2} \mathbf{grad}^2 \mathbf{tr} \right] \varphi = 0 = \mathbf{tr}^2 \varphi. \quad (6)$$

For example, if the symmetric form φ is a vector these are Maxwell's equations *in vacua*. When $\varphi = h_{\mu\nu} dx^\mu dx^\nu$ —metric fluctuations—the above equations are the linearized Einstein equations. Using the superalgebra (3,4), our Kähler equations mimic these massless higher spin equations

$$\left[\Delta - 2\partial\bar{\partial} - 2\bar{\partial}\partial + 2\partial\bar{\partial}\Lambda \right] A = 0. \quad (7)$$

¹Supersymmetric mechanics on Kähler manifolds has been extensively studied in [6].

where $(\partial, \bar{\partial})$ play the *rôle* of **grad**, $(\delta, \bar{\delta})$ of **div** and the Kähler form contraction Λ of **tr**. For scalars, the equations (7) reduce to the Laplace equation. In that case, in four real dimensions, viewing $F = \partial\bar{\partial}\varphi$ as an abelian field strength, the resulting equation $\Lambda F = 0$ is the anti-self-duality condition of [7]. For a $(1,0)$ -form $A = A_i dz^i$ we obtain simply $\delta\partial A = 0$, the natural complex generalization of the Maxwell system. *I.e.*, for vectors we obtain a holomorphic and anti-holomorphic copy of Maxwell's equations². Note, our system is not equivalent to the standard equations $d^*dA = 0$.

Our Kähler equations clearly enjoy gauge invariances

$$A \sim A + \partial\alpha + \bar{\partial}\bar{\alpha} \quad (8)$$

for arbitrary forms α and $\bar{\alpha}$. A corresponding pair of Bianchi identities hold thank to the operator facts

$$[\delta - \frac{1}{2}\partial\Lambda] \wedge \partial\bar{\partial} = 0 = [\bar{\delta} + \frac{1}{2}\bar{\partial}\Lambda] \wedge \partial\bar{\partial}. \quad (9)$$

These Bianchi identities seem to bear little relation to the gauge invariances (8). In fact, since we claim that our equations are the natural generalization of Maxwell's equations $d^*dA = 0$, the apparent asymmetry of our equations under adjoints seems disappointing. Moreover, if A is a general form, we also expect gauge for gauge symmetries analogous to those for higher-form electromagnetism. In fact both these objections are easily remedied.

Firstly, if we multiply the Kähler equations³ on the left by the operator

$$: I_1(2\sqrt{L\Lambda})/\sqrt{L\Lambda} :$$

where $:\bullet:$ denotes normal ordering by form degree (*i.e.* with L to the left and Λ the right) and the modified Bessel function of the second kind obeys the asymptotic expansion⁴ $I_1(2\sqrt{z})/\sqrt{z} = 1 + \frac{1}{2}z + \frac{1}{12}z^2 + \frac{1}{144}z^3 + \frac{1}{2880}z^4 + \dots$, then we find the symmetric form for our equations

$$GA = 0 \quad (10)$$

where the self-adjoint operator G is given by

$$G = : \frac{I_0(2\sqrt{L\Lambda})}{\sqrt{L\Lambda}} (\Delta - 2\partial\bar{\delta} - 2\bar{\partial}\delta) + 2 \frac{I_1(2\sqrt{L\Lambda})}{\sqrt{L\Lambda}} (\partial\bar{\partial} + \delta\bar{\delta}) : , \quad (11)$$

²Holomorphic topological Yang–Mills theory was studied in [8, 9].

³An analogous trick makes the symmetric tensor equations (6) self-adjoint.

⁴This series terminates in any given dimension when expanded in terms of Λ .

where all operators are normal ordered by form degree. In turn, a gauge invariant action principle is $S = \int_M (A, GA)$.

We complete this Note by explicating the gauge for gauge symmetries of our system. In the above symmetric formulation the gauge invariances and Bianchi identities are encapsulated by the operator identities

$$\delta G = \bar{\delta} G = G\partial = G\bar{\partial} = 0. \quad (12)$$

If we introduce auxiliary, commuting variables $(p, \bar{p})$ and the nilpotent operator

$$D = \partial \frac{\partial}{\partial p} + \bar{\partial} \frac{\partial}{\partial \bar{p}}, \quad (13)$$

then we can rewrite the gauge invariance (8) as $A \sim A + Da$ where a is any form linear in the auxiliary variables $(p, \bar{p})$. However, since D squares to zero on any (analytic) function of $(p, \bar{p})$ we obtain a complex

$$\dots \xrightarrow{D} \mathcal{F} \xrightarrow{D} \mathcal{F} \xrightarrow{D} \mathcal{F} \xrightarrow{D} \dots,$$

where $\mathcal{F}$ denotes forms depending on $(p, \bar{p})$. This complex precisely encodes the gauge for gauge symmetries of our Kähler equations.

The higher form Kähler electrodynamics announced in this Note were originally constructed by BRST quantization of a Kähler spinning particle model (see [4] for early studies). Similar Kähler systems and related higher form equations of motion involving half of the SUSY generators have been constructed in [5]. A detailed description of these techniques will appear in a forthcoming paper [10]. We thank Fiorenzo Bastianelli, Stefano Bellucci, Roberto Bonezzi, Ori Ganor, Andy Neitzke, Boris Pioline, Martin Roček, Stanley Deser and Bruno Zumino for stimulating discussions.

References

- [1] P. Griffiths and J. Harris, “Principles of algebraic geometry”, Wiley 1978.
- [2] J. M. Figueroa-O’Farrill, C. Kohl and B. J. Spence, Nucl. Phys. B **503**, 614 (1997) [arXiv:hep-th/9705161].
- [3] T. Damour and S. Deser, Annales Poincare Phys. Theor. **47**, 277 (1987); K. Hallowell and A. Waldron, Nucl. Phys. B

- 724** (2005) 453 [arXiv:hep-th/0505255], SIGMA **3**, 089 (2007), arXiv:0707.3164 [math.DG], Commun. Math. Phys. **278** (2008) 775 [arXiv:hep-th/0702033]; A. Lichnerowicz, Inst. Hautes Etudes Sci. Publ. Math. **10** (1961); Bull. Soc. Math. France **92**, 11 (1964); J. M. F. Labastida, Nucl. Phys. B **322**, 185 (1989); M. A. Vasiliev, Phys. Lett. B **209**, 491 (1988).
- [4] N. Marcus and S. Yankielowicz, Nucl. Phys. **B** 432 (1994) 225 [arXiv:hep-th/9408116], N. Marcus, Nucl. Phys. **B** 439, 583 (1995) [arXiv:hep-th/9409175].
- [5] F. Bastianelli and R. Bonezzi, “ $U(N)$ spinning particles and higher spin equations on complex manifolds”, arXiv:0901.2311[hep-th].
- [6] S. Bellucci and A. Nersessian, Phys. Rev. D **64** (2001) 021702 [arXiv:hep-th/0101065]; S. Bellucci and A. Nersessian, Nucl. Phys. Proc. Suppl. **102** (2001) 227 [arXiv:hep-th/0103005]; S. Bellucci, S. Krivonos and A. Nersessian, Phys. Lett. B **605** (2005) 181 [arXiv:hep-th/0410029].
- [7] M. Itoh, Publ. Res. Inst. Math. Sci. Kyoto **19**, 15 (1983).
- [8] A. Galperin and O. Ogievetsky, Commun. Math. Phys. **139**, 377 (1991).
- [9] J. S. Park, Nucl. Phys. B **423**, 559 (1994) [arXiv:hep-th/9305095].
- [10] D. Cherney, E. Latini and A. Waldron, in preparation.